

IF YOU ARE IN
TREATMENT,
DO NOT STOP TAKING
YOUR MEDICATION AT
THE FIRST SIGNS OF
RECOVERY.

COMPLETE THE TREATMENT.



Consult the nearest healthcare facility if you have symptoms such as **expectoration, fever, night sweats, weight loss, loss of appetite, and fatigue accompanied by two-plus weeks of coughing.**

*This publication was produced with the financial support of the European Union.
Its contents are the sole responsibility of the Republic of Türkiye Ministry of Health and do not necessarily reflect the views of the European Union.*

RIGHT PLACE, QUALITY SERVICES

Contact the nearest **Migrant Health Centre** for more information and support



www.sihhatproject.org



[sihhatproject](https://www.youtube.com/sihhatproject)




This project is funded by the European Union.
Bu proje Avrupa Birliği tarafından finanse edilmektedir.
هذا المشروع تم تمويله من قبل الاتحاد الأوروبي

**WE AIM FOR A
TUBERCULOSIS-
FREE TÜRKİYE**



**TREATMENT
FOR
TUBERCULOSIS
(PHTHISIS)
IS FREE OF
CHARGE IN
TÜRKİYE.**



Tuberculosis is an infectious disease that can be treated.

TB treatment goes on for at least 6 months. Treatment may take longer for drug-resistant cases (18-24 months).

For the majority of patients, infectiousness clears 2-3 weeks after starting the treatment.

TB is treatable with a regularly followed and completed medication course, which also prevents the spread of tuberculosis.



Directly Observed Treatment;

A tuberculosis patient takes each dose of medication in the presence of an observer during the entire treatment.

Directly Observed Treatment is essential and effective for effective treatment.

Do not stop taking your medication when you see signs of recovery after starting treatment. Complete the treatment.

Irregular use of medication makes the treatment more difficult causing infectiousness to continue.

Irregular use of medication leads to resistant cases of tuberculosis.

Resistant tuberculosis requires more medication to be taken for longer.

Contact your physician as soon as possible if you experience an unexpected symptom while taking your medication.

